



Published in final edited form as:

Clin Ther. 2013 January ; 35(1): A18–A32. doi:10.1016/j.clinthera.2012.12.014.

Childhood Obesity: A Review of Increased Risk for Physical and Psychological Co-morbidities

Elizabeth R. Pulgarón, PhD¹

¹University of Miami, Miller School of Medicine, Department of Pediatrics, Division of Clinical Psychology, P.O. Box 016820 (D-820), Miami, FL 33101

Abstract

Background—Worldwide estimates of childhood overweight/obesity are as high as 43 million and rates continue to increase each year. Researchers have taken interest in the childhood obesity epidemic and the impact of this condition across health domains. The consequences of childhood and adolescent obesity are extensive, including both medical and psychosocial comorbidities.

Objective—The purpose of this review was to consolidate and highlight the recent literature on the comorbidities associated with childhood obesity, both nationally and internationally.

Methods—PubMed and PsychINFO searches were conducted on childhood obesity and comorbidities.

Results—The initial search of the terms “obesity” and “comorbidity” yielded over 5000 published articles. Limits were set to include studies on children and adolescents that were published in peer-reviewed journals from 2002 to 2012. These limits narrowed the search to 938. Review of those articles resulted in 79 that are included in this review. The major medical comorbidities associated with childhood obesity in the current literature are metabolic risk factors, asthma, and dental health issues. Major psychological comorbidities include internalizing and externalizing disorders, ADHD, and sleep problems.

Conclusions—The high prevalence rates of childhood obesity have resulted in extensive research in this area. Limitations to the current childhood obesity literature include differential definitions of weight status and cut off levels for metabolic risk factors across studies. Additionally, some results are based on self-report of diagnoses rather than chart reviews or physician diagnosis. Even so, there is substantial support for metabolic risk factors, internalizing disorders, ADHD, and decreased health related quality of life as comorbidities to obesity in childhood. Additional investigations on other diseases and conditions that may be associated with childhood obesity are warranted and intervention research in this area is critical.

Keywords

Childhood Obesity; Medical Co-morbidities; Psychological Co-morbidities

© 2013 Excerpta Medica, Inc. All rights reserved.

All correspondence concerning this article should be addressed to Elizabeth R. Pulgaron, PhD., Department of Pediatrics, Division of Clinical Psychology, P.O. Box 016820 (D-820), Miami, FL 33101. epulgaron@med.miami.edu. Telephone: 305-243-0807 Fax: 305-243-4512.

“Conflict of interest statement”: The author has no conflict of interests to declare.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Obesity is the result of a chronic caloric imbalance, with more calories being consumed than expended each day. History of obesity, hereditary factors, environment, metabolism, behavior, culture, and socioeconomic status all play a role in obesity.¹ Most obese adults were obese as adolescents and most obese adolescents were overweight and/or obese as children.² In fact, the origins of obesity are being traced to early childhood development. Children who experience early adiposity rebound (before the age of five years) have increases in mean body mass index (BMI) from age three to adolescence while those that experience late adiposity rebound have decreases in BMI from age three to adolescence; differences between those who experience early and late adiposity rebound are maintained into adulthood.^{3,4} Evidence is available to support both genetic^{5,6} and environmental components^{7,8} to obesity. Availability of healthy snacks and meals⁹ and children's choices in food consumption¹⁰ are also important to consider.

In addition to consumption of food, energy expenditure, generally in the form of physical activity, is highly important to maintaining healthy weight. Results from the 2007 Youth Risk Behavioral Survey conducted in the US indicated that among high school students nationwide, 35% had watched television 3 or more hours per day on an average school day during the past month and 65% had not met recommended levels of physical activity during the past week.¹¹ Recent estimates from the National Health and Nutrition Examination Survey¹² indicate that approximately one third of children in the United States are overweight or obese, with approximately 17% meeting criteria for obesity as measured by a BMI score at or above the 95th percentile.¹³ BMI scores can be converted into standardized scores, zBMI, to conduct comparisons between groups.¹⁴ Ethnic minority children are at an increased risk for obesity.¹⁵ Worldwide estimates of childhood overweight/obesity are as high as 43 million. This number reflects an increase from 4.2% in 1990 to 6.7% in 2010.¹⁶

The rates of obesity in youth are expected to continue to grow. As a result it is imperative to consider the physical health and psychological correlates and consequences of this condition. The consequences of childhood and adolescent obesity are far reaching, not only including health-related physical outcomes, such as high blood pressure, high cholesterol, metabolic syndrome, type 2 diabetes, orthopedic problems, sleep apnea, asthma, and fatty liver disease, but also psychological, social and behavioral consequences, such as risk for problems related to body image, self-esteem, social isolation and discrimination, depression, and reduced quality of life.¹ In 2003 Reilly and colleagues¹⁷ conducted a systematic review of the literature on the health consequences of childhood obesity. They concluded that childhood obesity has significant short term and long-term adverse medical and psychosocial effects extending into adulthood. Strong evidence exists for childhood obesity affecting morbidity and mortality in adulthood.¹⁸ The purpose of this review is to explore and expand upon previously identified associations between childhood obesity with physical and psychological co-morbidities.

Methods

Electronic searches were conducted via PubMed and PsychInfo in October of 2012 using the search terms "obesity" and "comorbidity". The following types of publications were excluded from this review: letters to editors, commentaries, case studies, and review articles. All other publications were considered for inclusion. Articles that did not provide data on medical or psychological co-morbidities of obesity were excluded. Clinical trials and randomized control trials that included baseline information on prevalence of comorbidities in obese children compared to a healthy weight control group were included, otherwise trials were excluded. Most of the studies included in this review were cross sectional, including

one time assessments, retrospective chart reviews, and national database analyses. Some studies were prospective or analyzed trends across time. Although the intended sample for this review was pediatric (under the age of 18 years-old), some studies included participants up to the age of 22. Age limits were extended to allow for these studies to be included in the review.

Results

The PubMed search initially yielded 4315 results. Limits were applied to narrow the search including being published from 2002 to 2012 (3800 results), age range of participants was between 0 and 18 years old (961), written in English (840 results). A similar search was conducted in Psych Info. This search initially yielded 709 results, using the same limits described above the search was narrowed to 53 articles. Combining the two searches resulted in 893 articles. The author and a research assistant read all abstracts to further narrow the search to 291 articles for in depth review. This list was narrowed down to 79 articles that focused on comorbidities associated with childhood obesity. The 79 articles that comprise this review were organized into four tables based on type of outcome variable (medical or psychological) and country of research (US or international). If medical and psychological outcomes were assessed in one study, then it was listed in both tables. See tables 1 and 2 for details on variable of interest, study population descriptive, and author details.

Medical Co-morbidities

The main medical co-morbidities associated with childhood obesity throughout this review include asthma, metabolic risk factors, and dental health. Some evidence was also available to support a link between childhood obesity and other medical conditions.

Asthma—The majority of the literature available on obesity and asthma in children indicates that there is a relationship between the two conditions^{19–21} especially when symptoms of asthma versus an official diagnosis is studied.^{22–26} Even in areas where rates of obesity are very low, there are still some associations between obesity and asthma (adjusted OR = 2.36; 95 % CI, 1.02 – 5.44; $P=0.04$; $n=2926$).²⁷ Studies have found differential effects between genders, with some supporting a relationship between obesity and asthma for boys but not for girls (adjusted OR = 2.36; 95 % CI, 1.02–5.44; $P=0.04$; $n=2926$)^{19,27} and others vice versa (OR = 2.73; 95% CI = 1.09 – 6.85, $P=0.032$ (girls); OR = 1.74; 95% CI, 0.83–3.73, $P=0.137$ (boys); $n=854$).^{22,26}

However, there are studies that do not document an association between obesity and asthma, particularly in minority populations²⁶ or abroad.^{28,29} For instance, Vignolo and colleagues²⁹ did not find differences in rates of asthma between children who were obese and those who were not ($P=.08$, $n=1179$) in a sample of Italian youth. Authors propose differences in diet across countries as a potential explanation. Other researchers recognize that there is a relationship between obesity and asthma, but are unable to verify whether increases in obesity help explain the increases in asthma diagnoses. For example, Wickens and colleagues³⁰ found large increases in the prevalence of asthma diagnoses and medication use in New Zealand across a 10 year span, but statistical tests that adjusted for multiple comparisons indicated that increase in prevalence was not accounted for by increases in BMI ($P=.04$). There is also some evidence for the combination of obesity and asthma to predispose children to higher rates of metabolic risk factors.³³ For asthma related conditions such as other atopic diseases or allergies, results are mixed.^{25,34} Some researchers have found evidence to support a link between obesity or percent body fat and these conditions

(OR = 1.03; 95% CI, 1.01 – 1.06; $P = .017$)³⁵ and others have not (OR = .65, CI, .35 – 1.24, $P = .19$, $n = 2926$ of having eczema).^{25,27}

Metabolic Risk Factors—Although the exact criteria for metabolic syndrome differs across studies^{36,37}, the primary factors are large waist circumference, hypertension, high triglycerides, hyperglycemia, and low HDL cholesterol. All 35 studies included in this review that examined some aspect of metabolic risk found that overweight/obese children and adolescents are more likely to experience these risk factors than their healthy weight peers^{38–40} (see Tables 1 and 2). For instance, in a sample of 186 obese youth from Turkey, nearly 80% had at least one cardiometabolic risk factor, many had more than one risk factor⁴¹. In a study of overweight or obese Australian youth ($n=107$), compared to healthy weight matched controls ($n=182$), youth who were overweight/obese experienced significantly higher levels of hypertension (chi square p value = .012), impaired glucose tolerance ($P = .037$), hyperinsulinism ($P < .001$), and raised alanine transaminase ($P < .001$).⁴² In a study of school aged Hispanic children who were followed for two years it was found that for every one unit increase in zBMI the odds ratio of meeting criteria for metabolic syndrome was 2.4 (CI, 1.21 – 4.63, $P < .01$).⁴³

High blood pressure is consistently related to obesity^{44–49} and one of the metabolic risk factors most seen in children. For example, a national study on 1,021,211 17-year-old adolescents from Israel, who were receiving a medical evaluation for military service documented increases in both BMI ($P < .0001$) and high blood pressure ($P < .001$) across a 14 year period. BMI was associated with high blood pressure (OR = 4.11, CI = 3.89–4.34 and OR = 5.56; CI, 5.09–6.07 for male and female subjects, respectively), type 2 diabetes (OR = 5.56; CI, 5.09–6.07 and OR = 4.42; CI, 3.90 – 5.00, for male and female subjects, respectively) and hyperlipidemia (OR = 16.07, CI, 8.29 – 31.15 and OR = 9.00; CI, 4.36–18.6 for male and female subjects, respectively) even when controlling for origin, level of education and the year of recruitment. Severe hypertension was associated with BMI for males ($P < .0001$), but not females ($P = .053$).⁵⁰ Similar trends regarding high blood pressure and obesity have been reported in American youth, especially in the last 10 years.⁵¹ Obese youth are twice as likely to have hypertension (for SBP > 140, OR = 2.24; CI, 1.46 – 3.45; $P < 0.001$, and for DBP > 90, OR = 2.10; CI, 1.063–4.17, $P = 0.03$)⁵² and high blood pressure has been documented as a co-morbidity of obesity in minority and immigrant samples as well.^{28,45,53} One study of 1053 patients from an obesity outpatient clinic in Germany found that rates of metabolic syndrome were significantly higher among Turkish patients (40.4%) compared to Germans (27.3%; $p = 0.02$).⁵³

Many studies demonstrated that the degree of obesity impacts level of metabolic risk, with those in the extreme obesity range experiencing the worst outcomes.^{37,54} For instance, among adolescents with a BMI above the 97th percentile for their age Lafortuna and colleagues documented rates of metabolic syndrome as high as 40.4% for the German sample and 23.3% for the Italian sample.³⁷ Evidence for the high rates of metabolic risk factors have been documented in both clinic and population-based settings in extremely obese ethnic minority populations and subgroups.⁵⁵

Dental Health—The role of childhood obesity on dental health has become more prevalent in recent years, with research in this area being conducted with children as young as two. Associations between obesity and dental health have been assessed nationally and internationally. For example, Willerhausen and colleagues⁵⁶ documented a significant positive correlation ($P = .002$) between BMI and dental health (caries frequency) in a group of 2071 elementary school aged children in Germany. Similar variables were assessed in a national database study conducted in the US with younger children (ages 2–6).⁵⁷ The association between childhood obesity and caries was not significant after controlling for

age, race, and poverty/income ratio. However, in the oldest subgroup of the study, the 60–72 month age group, the relationship was significant ($P = .049$). This age group is closer to the lower end of the age range for the Willershausen study⁵⁶ indicating that perhaps the relationship between obesity and caries is present for children at certain ages. The relationship between obesity and caries has been documented in other studies as well.^{58,59} Marshall and colleagues⁶⁰ found a relationship between children who are at risk for overweight ($P < .05$), (but not for overweight) and caries. McGuire and colleagues⁶¹ investigated another aspect of dental health, erosive tooth wear, but found no differences between healthy weight and overweight children.

Other Potential Medical Co-morbidities—In addition to the more established co-morbidities described above, there are a number of additional medical conditions that have been associated with childhood obesity in the literature. For instance, four studies investigated the potential role of obesity on non-alcoholic fatty disease (NAFD). With some supporting the role of obesity in NAFD^{69–71} and others not.⁷² One example of a study which provided evidence of the link between obesity and NAFD was conducted by Adibi and colleagues. This cross-sectional study was conducted with healthy children between the ages of 6 to 18 years from Isfahan schools, who had been randomly selected for the Isfahan Healthy Heart Program. They found that the prevalence of sonographic fatty liver (SFL) in obese children was 54.4%, which was significantly higher the percent of SFL found in overweight (10.5%) and healthy weight children (1%; $P < 0.001$). Two studies ascertained an association between obesity and gastrointestinal (GI) problems. In a study of 1156 children Stordal and colleagues found that overweight children were nearly twice as likely to report GERD than healthy weight peers (OR = 1.8; 95% CI, 1.2–2.6).⁷³ In the second study which compared obesity rates of 757 patients from a GI clinic and 255 matched controls results indicated significantly higher rates of obesity for the GI sample, $P < .001$.^{73,74}

Some studies have found that children and/or adolescents who are obese are also more likely to have back pains, iron deficiency, disc degeneration, foot problems, elevated serum thyroid-stimulating hormone (TSH), endothelial dysfunction in children, acanthosis nigricans, increased carotid intima media thickness (IMT), differences in timing of sexual maturation, and use antibiotics^{20,75–83} but not kidney stone disease.⁸⁴

Psychological Comorbidities

In addition to the traditional medical co-morbidities associated with childhood obesity, researchers are investigating the psychosocial sequelae that accompany obesity at a young age. Research in this area focus on ADHD, internalizing and externalizing disorders, and sleep. See tables 3 and 4 for details.

Attention Deficit Hyperactivity Disorder (ADHD)—A number of studies have assessed the relationship between obesity and ADHD and many researchers suggest that the dysregulation these children experience with their behavior extends into their eating habits. Diagnoses of ADHD have been found to be more prevalent in obese children than their healthy weight peers.^{85,86} For instance, Erhart and colleagues⁸⁷ found that after controlling for age, gender, and socio-economic status, overweight/obese children were found to be twice as likely to have an ADHD diagnosis (OR = 2.0; 95% CI, 1.23 – 3.11). Not all studies have found an association between obesity and ADHD.²⁸ For example, a large scale community study conducted by Rojo and colleagues of 35,403 13–15 year olds did not support higher rates of ADHD in obese samples.⁸⁸ In fact, analyses indicated lower rates of ADHD in the obese group ($X^2 = 19.1$, $P < 0.001$; (OR) = 1.24; 95 (CI), 1.13–1.39) compared to healthy and underweight peers. The main limitation of the Rojo study⁸⁸ is that ADHD was assessed with a self-report measure and not through a standardized clinical

assessment. The self-reported rates of ADHD in the sample were much higher (24%) than typical prevalence rates (4%). Additionally, Marks and colleagues⁸⁹ reported that in a sample of 40 children with ADHD seeking therapy through a telepsychiatry clinic were actually more likely to be underweight or normal weight than overweight or obese ($t(106.379) = 1.991, P = 0.049$).

Internalizing/Externalizing Disorders—Higher rates of internalizing and externalizing symptoms and/or disorders have been documented in obese youth compared to healthy weight peers^{21,42,90–96}. In comparison to a sample of children with diabetes, children who were obese displayed significantly higher internalizing (50 out of 116 cases vs. 40 out of 155 cases; $p = .004$) and externalizing symptoms (29 out of 116 vs. 19 out of 155; $P = .010$) on parent report questionnaires.⁹⁷ However, analyses within the obese group indicated that those in the “super-obese” group (defined by a BMI z-score ≥ 8) had lower symptoms of anxiety Student’s t test: ($t = 2.795, P = .006$) and depression ($t = 2.180, P = .031$) compared to other obese children. In some studies the relationship between obesity and depression is not consistent across subgroups. For example, after adjusting for gender, age, race/ethnicity and poverty status a link between obesity and depression was documented in males (adjOR = 2.7; 95% CI, 1.1–7.1) and non-Hispanic Blacks (adjOR = 3.1; 95% CI, 1.1–8.3), but not for the overall group of 4,150 adolescents (adjusted odds ratio (adjOR) = 1.6; 95% CI, 0.9 – 2.9) who were included in a study using data from the National Health and Nutrition Examination Survey.⁹⁸ In a prospective study Anderson and colleagues⁹⁵ found that the presence of internalizing and externalizing disorders in females, but not males, was associated with increases in zBMI, .13 and .09, $P < .05$, respectively. Marks and colleagues⁸⁹ did not find differences on depression between individuals who were obese and those who were not in a smaller study of 31 patients receiving treatment through a telepsychiatry clinic. Other researchers who may not have documented a relationship between anxiety and depression and weight status, have found significant relationships with similar variables and weight status. For instance in a sample of 8,090 students from Norway poor self-esteem, but not anxiety and depression, was associated with overweight and obesity in male (OR = 0.9; CI, 0.86–0.94) and female (OR = 0.8; CI, 0.74–0.92) adolescents.⁹⁹ Data from large school-based cohort ($n = 10\,403$) in China indicated that the perception of weight status, but not BMI, was related to psychopathology (OR = 1.61; CI, 1.17 – 2.20 for males; OR = 1.88; CI, 1.54 – 2.31 for females).¹⁰⁰

The extent of behavioral problems in overweight children can be quite severe. For instance, in one study using data from a nationally representative sample of kindergartners in the United States ($n = 9949$) overweight girls had over 81% greater odds of having substantial teacher-reported externalizing behavior problems compared with girls who were not overweight, OR = 1.81; 95% CI, 1.23 – 2.68.⁹⁴ As with many of the associated comorbidities of childhood obesity, it is difficult to determine the temporal relationship between obesity and behavioral problems. Many propose behavioral problems are a result of the stigmatization associated with childhood obesity, but there is also evidence to support that behavioral problems may precede overweight status in some children.¹⁰¹ In a sample of 629 youth clinically significant behavior problems in normal-weight children were independently associated with becoming overweight 2 years later (adjusted OR = 5.23; 95% CI, 1.37–19.93) after adjusting for covariates.

Sleep—Various researchers have examined the role of obesity on sleep. Studies indicate obesity is associated with short sleep duration.^{108,109} Specifically, one study showed that compared with healthy weight peers, overweight children slept about 22 minutes less on average ($\beta = -0.174; P = .02$), had lower sleep efficiency ($\beta = -0.027; P = .01$), lower REM density ($\beta = -0.256, P = .02$).¹¹⁰ In addition to the duration and quality of sleep, the presence of obstructive sleep apnea syndrome (OSAS) in obese children is concerning.

Studies indicate that OSAS is more frequent¹¹¹ and severe¹¹² in overweight/obese children compared to healthy weight controls. Kohler and colleagues¹¹³ documented a 3.5 time increase for each standard deviation increase in zBMI among adolescents, OR = 3.55; 95% CI, 1.30 – 9.71; $P = .01$. On the other hand, OSAS was not associated with BMI in a clinic sample of Hispanic youth²⁸ and sleep problems were not significantly higher in obese youth compared to healthy weight peers in an Australian sample.^{49,114}

Discussion

The purpose of this review was to summarize the current literature on the medical and psychological comorbidities associated with childhood obesity across national and international investigations. Overall, there are certain medical and psychological comorbidities associated with childhood obesity that have been fairly well established in the literature.

Medically, the association between childhood obesity and metabolic risk factors is evident.^{52,115–117} Other medical comorbidities, such as asthma, have received significant research and fairly consistently a relationship between the two conditions is reported,^{19–21} but there are exceptions in certain subsamples.^{29,30} Researchers have suggested differences in diet across cultures as a potential explanation for the lack of relationship between obesity and asthma in certain subgroups.²⁹ As with many potential co-morbidities, the causal link between obesity and asthma remains uncertain since many children with asthma avoid physical activity to reduce likelihood of experiencing symptoms resulting in less energy expenditure and more likelihood of gaining weight.³¹ It still needs to be determined whether healthy weight helps regulate asthma symptoms and/or whether good asthma control helps maintain a healthy weight.³² In general, there is some support for an association between obesity and dental problems,^{56,58,59} but more research is needed in the area to decipher the role of potential mediators, such as poor dietary habits and age, that might explain the cause of dental problems in children. Many other medical comorbidities have been associated with childhood obesity in some regard^{75–83,118} but insufficient evidence has been presented to conclude that there is a clear relationship between the two. The overlap between medical co-morbidities is another difficulty researchers encounter in this line of research is. Many of the published studies control statistically for the effect of age, gender, SES and other demographic factors that could be related to co-morbidities, however, there are so many potential confounds and so much interdependency among the co-morbidities that it is difficult for researchers to isolate the effects of childhood obesity. One confound that could be better accounted for in the literature is the effect of puberty. Controlling for age does not suffice for individual differences in rates of development. Although some studies account for pubertal status,⁸² others do not.^{119,120} Pubertal status can be very influential, especially for medical comorbidities.

Within the psychological domain there are many studies which propose that obesity is associated with increased internalizing and externalizing symptoms, ADHD diagnoses, and sleep problems^{85,94,95,108}, but these relationships are not certain because other studies have not been able to establish these associations or have found inverse relationships.^{88,89} One of the greatest challenges within the psychological domain is the definition of the psychological variable of interest. Participants are often asked to self-report diagnoses rather than having chart reviews conducted, or ideally clinician administered structured interviews or assessments. These methods may be a function of cost and feasibility, especially for large-scale longitudinal trials, but they must be considered when interpreting results. Another potential explanation for the lack of consistency in psychological co-morbidities of obesity is the often underestimated value of resiliency. There is significant variation between levels of depression and anxiety documented among studies of obese children. It is likely

that other variables, such as level of self-esteem or family support, may explain the relationship (or lack there of) in certain groups. More sophisticated and complex psychosocial modeling will be necessary to help explain the relationship between obesity and internalizing symptoms.

Part of the challenge with this topic area is that there are so many potential comorbidities of obesity that the literature seems oversaturated in some ways, but understudied in others. Across both medical and psychological domains the conclusions that can be drawn from the current literature are limited by the methodology through which data were collected. Although some studies are prospective or at least contain a prospective element within the study design,^{49,65} most of the studies are cross sectional and therefore the direction of the relationships cannot be established. In many cases it appears that obesity occurs first and then the subsequent comorbidity arises, but causality cannot be determined until more prospective studies are conducted. Part of the challenge with the call for prospective studies is the length of time between a child being diagnosed as obese and the appearance of comorbidities, with some co-morbidities not being diagnosed until later adulthood. Another consideration with prospective studies is how young children are becoming obese. It is well established in the literature that obesity rates in preschoolers are increasing rapidly¹⁶, yet only 4 of the articles in this review focused on children under the age of 5. In order to obtain a more comprehensive understanding of the co-morbidities associated with childhood obesity, researchers will need to focus investigations on younger populations.

The definition of childhood obesity, a basic, yet fundamental concept to this area of research is a consistent challenge encountered by researchers in this domain. Many studies conducted in the US in the past 10 years adhere to the same age and gender zBMI percentiles recommended by the Center for Disease Control 85th percentile for overweight and 95th percentile for obese.¹²¹ Others use 90th percentile for overweight and 97th percentile for obese.⁸⁷ The World Health Organization uses 2 standard deviations above the WHO growth standard median as guidelines for overweight and does not use the term obese.¹²² Other researchers use the International Obesity Task Force guidelines¹²³ or guidelines specific to the country the research is being conducted.⁵³ Some investigators use adult overweight (BMI>25) and obesity (BMI>30) guidelines.⁷³ Although the merits of each of these classification systems can be debated, the lack of consistency among BMI thresholds used by researchers makes consolidating evidence across studies problematic and a limitation of the literature as a whole. BMI also needs to be assessed more rigorously. Although the majority of studies used trained staff to administer anthropometric assessments, there are studies that use self-reported weight and height, a limitation of some of the large national database studies.¹²⁴ In addition to variations in the definition and measurement of obesity, there are also inconsistencies regarding the cut off points for the metabolic risk factors, which makes it very difficult to compare findings across studies and identify trends.³⁷ It is important to set consistent criteria for MS in children and adolescents in order for researchers' efforts to be maximized and not duplicated. It is possible that some of the inconsistencies in findings across studies are due to variations in definitions of key terms such as obesity and MS.

Seventy-nine studies were identified in a 10 year time span on the topic of childhood obesity and co-morbidities and the number being published continues to grow. Despite the recognition of the importance of this condition in children, the rates of childhood obesity continue to increase and unfortunately the reality of the long-term ramifications is becoming solidified in the literature. There is evidence to support significant medical and psychological sequelae of the condition into adulthood. The current literature supports the importance of continued work in this area to further understand the complexities of the intertwined comorbidities associated with childhood obesity. Future research is needed to

determine a more precise course for children who are obese and the need to continue intervention efforts in this population cannot be understated.

Acknowledgments

The author would like to thank the Pediatric Research Team at the University of Miami for their technical support in preparing this manuscript, especially Ashley Marchante for her assistance with reviewing the literature. The author would also like to recognize Drs. Alan Delamater, John Elder, and Sarah Messiah for their guidance on this manuscript. This work was supported in part by NHLBI grant number 3R01HL102130-02S1.

References

1. Delamater, AM.; Pulgaron, ER.; Daigre; O'Donohue, A.; Benuto, L.; Tolle, L., editors. Adolescent health psychology. New York, New York: Springer; 2013. Forthcoming
2. Rooney BBL. Predictors of obesity in childhood, adolescence, and adulthood in a birth cohort. *Matern Child Health J.* 2011; 15(8):1166–1175. [PubMed: 20927643]
3. Rolland-Cachera MF, Deheeger M, Maillot M, Bellisle F. Early adiposity rebound: Causes and consequences for obesity in children and adults. *Int J Obes.* 2006; 30:S11–S17.
4. Williams SMGA. Patterns of growth associated with the timing of adiposity rebound. *Obesity.* 2009; 17:335–341. [PubMed: 19057527]
5. Allison, DB.; Matz, PE.; Pietrobelli, A.; Zannolli, R.; Faith, MS. Genetic and environmental influences on obesity. In: Bendich, A.; Deckelbaum, RJ., editors. Primary and secondary preventive nutrition. Totowa, NJ: Humana Press; 1999. p. 147-164.
6. Meyer, JMSA. Genetics and human obesity. In: Stunkard, AJ.; Wadden, TA., editors. *Obesity: Theory and therapy.* 2. New York, NY: Raven Press; 1993. p. 137-149.
7. Kubik MY, Lytle LA, Story M. Schoolwide food practices are associated with body mass index in middle school students. *Arch Pediatr Adolesc Med.* 2005; 159:1111–1114. [PubMed: 16330732]
8. Miller J, Gold MS, Silverstein J. Pediatric overeating and obesity: An epidemic. *Psychiatric Annals.* 2003; 33(2):94–103.
9. Goslinger W, Madsen K, Woodward-Lopez G, Crawford P. Would students prefer to eat healthier foods at school? *J Sch Health.* 2011; 81(3):146–151. [PubMed: 21332479]
10. Parks S, Sappenfield W, Huang Y, Sherry B, Bensyl D. The impact of the availability of school vending machines on eating behavior during lunch: The youth physical activity and nutrition survey. *J Am Diet Assoc.* 2010; 110:1532–1536. [PubMed: 20869493]
11. Eaton D, Kann L, Kinchen S, Shanklin S, Ross J, Hawkins J, Wechsler H. Youth risk behavior surveillance—United states, 2007. *MMWR Surveill Summ.* 2008; 54(4):1–131. [PubMed: 18528314]
12. Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high body mass index in US children and adolescents, 2007–2008. *JAMA-J Am Med Assoc.* 2010; 303(3):242–249.
13. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA-J Am Med Assoc.* 2012; 307(5):483–490.
14. Kuczmarski RJ, Ogden CL, Guo SS, et al. CDC growth charts for the united states: Methods and development. *Vital Health Stat 11.* 2000; 2002(246):1–190.
15. Kumanyika S, Grier S. Targeting interventions for ethnic minority and low-income populations. *Future Child.* 2006; 16(1):187–207. [PubMed: 16532664]
16. de Onis M, Blossner M, Borghi E. Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr.* 2010; 92(5):1257–1264. [PubMed: 20861173]
17. Reilly JJ, Methven E, McDowell ZC, et al. Health consequences of obesity. *Arch Dis Child.* 2003; 88(9):748–752. [PubMed: 12937090]
18. Reilly JJJ. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: Systematic review. *International journal of obesity (2005).* 2011; 35(7):891–898. [PubMed: 20975725]
19. Gilliland FD, Berhane K, Islam T, et al. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J Epidemiol.* 2003; 158(5):406–415. [PubMed: 12936895]

20. Lazorick S, Peaker B, Perrin EM, et al. Prevention and treatment of childhood obesity: Care received by a state medicaid population. *Clin Pediatr (Phila)*. 2011; 50(9):816–826. [PubMed: 21525083]
21. Trent M, Jennings JM, Waterfield G, Lyman LM, Thomas H. Finding targets for obesity intervention in urban communities: School-based health centers and the interface with affected youth. *J Urban Health*. 2009; 86(4):571–583. [PubMed: 19472059]
22. Mahut B, Beydon N, Delclaux C. Overweight is not a comorbidity factor during childhood asthma: The GrowthOb study. *Eur Respir J*. 2012; 39(5):1120–1126. [PubMed: 21885396]
23. Jang AS, Lee JH, Park SW, Shin MY, Kim DJ, Park CS. Severe airway hyperresponsiveness in school-aged boys with a high body mass index. *Korean J Intern Med*. 2006; 21(1):10–14. [PubMed: 16646558]
24. Tai A, Volkmer R, Burton A. Association between asthma symptoms and obesity in preschool (4–5 year old) children. *J Asthma*. 2009; 46(4):362–365. [PubMed: 19484670]
25. Silva MJ, Ribeiro MC, Carvalho F, Goncalves Oliveira JM. Atopic disease and body mass index. *Allergol Immunopathol (Madr)*. 2007; 35(4):130–135. [PubMed: 17663921]
26. Spathopoulos D, Paraskakis E, Trypsianis G, et al. The effect of obesity on pulmonary lung function of school aged children in greece. *Pediatr Pulmonol*. 2009; 44(3):273–280. [PubMed: 19208374]
27. Vlaski E, Stavric K, Isjanovska R, Seckova L, Kimovska M. Overweight hypothesis in asthma and eczema in young adolescents. *Allergol Immunopathol (Madr)*. 2006; 34(5):199–205. [PubMed: 17064649]
28. Mirza NM, Kadow K, Palmer M, Solano H, Rosche C, Yanovski JA. Prevalence of overweight among inner city hispanic-american children and adolescents. *Obes Res*. 2004; 12(8):1298–1310. [PubMed: 15340113]
29. Vignolo M, Silvestri M, Parodi A, et al. Relationship between body mass index and asthma characteristics in a group of italian children and adolescents. *J Asthma*. 2005; 42(3):185–189. [PubMed: 15962875]
30. Wickens K, Barry D, Friezema A, et al. Obesity and asthma in 11–12 year old new zealand children in 1989 and 2000. *Thorax*. 2005; 60(1):7–12. [PubMed: 15618575]
31. Ulger Z, Demir E, Tanac R, et al. The effect of childhood obesity on respiratory function tests and airway hyperresponsiveness. *Turk J Pediatr*. 2006; 48(1):43–50. [PubMed: 16562785]
32. Abramson NW, Wamboldt FS, Mansell AL, Carter R, Federico MJ, Wamboldt MZ. Frequency and correlates of overweight status in adolescent asthma. *J Asthma*. 2008; 45(2):135–139. [PubMed: 18350405]
33. Del-Rio-Navarro BE, Castro-Rodriguez JA, Garibay Nieto N, et al. Higher metabolic syndrome in obese asthmatic compared to obese nonasthmatic adolescent males. *J Asthma*. 2010; 47(5):501–506. [PubMed: 20560825]
34. Irei AV, Takahashi K, Le DS, et al. Obesity is associated with increased risk of allergy in vietnamese adolescents. *Eur J Clin Nutr*. 2005; 59(4):571–577. [PubMed: 15702126]
35. Irei AV, Takahashi K, Le DS, et al. Obesity is associated with increased risk of allergy in vietnamese adolescents. *Eur J Clin Nutr*. 2005; 59(4):571–577. [PubMed: 15702126]
36. Golley RK, Magarey AM, Steinbeck KS, Baur LA, Daniels LA. Comparison of metabolic syndrome prevalence using six different definitions in overweight pre-pubertal children enrolled in a weight management study. *Int J Obes*. 2006; 30(5):853–860.
37. Lafortuna CL, Adorni F, Agosti F, et al. Prevalence of the metabolic syndrome among extremely obese adolescents in italy and germany. *Diabetes Res Clin Pract*. 2010; 88(1):14–21. [PubMed: 20096473]
38. Flechtner-Mors M, Thamm M, Wiegand S, et al. Comorbidities related to BMI category in children and adolescents: German/austrian/swiss obesity register APV compared to the german KiGGS study. *Horm Res Paediatr*. 2012; 77(1):19–26. [PubMed: 22104037]
39. Reinehr T, Wunsch R. Relationships between cardiovascular risk profile, ultrasonographic measurement of intra-abdominal adipose tissue, and waist circumference in obese children. *Clin Nutr*. 2010; 29(1):24–30. [PubMed: 19576664]

40. Ng KC, Lai SW. Application of anthropometric indices in childhood obesity. *South Med J*. 2004; 97(6):566–570. [PubMed: 15255423]
41. Serap S, Mevlut B, Inanc C, Ender S. Metabolic syndrome in childhood obesity. *Indian Pediatr*. 2007; 44(9):657–662. [PubMed: 17921554]
42. Bell LM, Curran JA, Byrne S, et al. High incidence of obesity co-morbidities in young children: A cross-sectional study. *J Paediatr Child Health*. 2011; 47(12):911–917. [PubMed: 21902753]
43. Patiño-Fernández, Anna Maria; Delamater, Alan M.; Sanders, Lee; Brito, Arturo; Goldberg, Ronald. A prospective study of weight and metabolic syndrome in young hispanic children. *Children's Health Care*. 2008; 37:316.
44. Krzyzaniak A, Kaczmarek M, Stawinska-Witoszynska B, Krzywinska-Wiewiorowska M. Prevalence of selected risk factors for cardiovascular diseases in adolescents with overweight and obesity. *Med Wieku Rozwoj*. 2011; 15(3):282–287. [PubMed: 22006474]
45. Mohan B, Kumar N, Aslam N, et al. Prevalence of sustained hypertension and obesity in urban and rural school going children in ludhiana. *Indian Heart J*. 2004; 56(4)
46. Glowinska B, Urban M, Koput A, Galar M. New atherosclerosis risk factors in obese, hypertensive and diabetic children and adolescents. *Atherosclerosis*. 2003; 167(2):275–286. [PubMed: 12818410]
47. Langens F, Dapper T, Nuboer R, van Weel C, van Binsbergen J. Co-morbidity obese children in family practice in the netherlands: The results of a pilot study. *Fam Pract*. 2008; 25 (Suppl 1):i75–8. [PubMed: 18826992]
48. Ribeiro JC, Guerra S, Oliveira J, Andersen LB, Duarte JA, Mota J. Body fatness and clustering of cardiovascular disease risk factors in portuguese children and adolescents. *Am J Hum Biol*. 2004; 16(5):556–562. [PubMed: 15368603]
49. Wake M, Canterford L, Patton GC, et al. Comorbidities of overweight/obesity experienced in adolescence: Longitudinal study. *Arch Dis Child*. 2010; 95(3):162–168. [PubMed: 19531529]
50. Levin A, Morad Y, Grotto I, Ravid M, Bar-Dayana Y. Weight disorders and associated morbidity among young adults in israel 1990–2003. *Pediatr Int*. 2010; 52(3):347–352. [PubMed: 19807878]
51. Din-Dzietham R, Liu Y, Bielo MV, Shamsa F. High blood pressure trends in children and adolescents in national surveys, 1963 to 2002. *Circulation*. 2007; 116(13):1488–1496. [PubMed: 17846287]
52. Movahed MR, Bates S, Strootman D, Sattur S. Obesity in adolescence is associated with left ventricular hypertrophy and hypertension. *Echocardiography*. 2011; 28(2):150–153. [PubMed: 21276070]
53. Dannemann A, Ernert A, Rucker P, et al. Ethnicity and comorbidities in an overweight and obese multiethnic childhood cohort in berlin. *Acta Paediatr*. 2011; 100(4):578–584. [PubMed: 21223371]
54. l'Allemand D, Wiegand S, Reinehr T, et al. Cardiovascular risk in 26,008 european overweight children as established by a multicenter database. *Obesity (Silver Spring)*. 2008; 16(7):1672–1679. [PubMed: 18451769]
55. Messiah SE, Carrillo-Iregui A, Garibay-Nieto G, Lopez-Mitnik G, Cossio S, Arheart KL. Inter- and intra-ethnic group comparison of metabolic syndrome components among morbidly obese adolescents. *J Clin Hypertens (Greenwich)*. 2010; 12(8):645–652. [PubMed: 20695945]
56. Willershausen B, Moschos D, Azrak B, Blettner M. Correlation between oral health and body mass index (BMI) in 2071 primary school pupils. *Eur J Med Res*. 2007; 12(7):295–299. [PubMed: 17933701]
57. Hong L, Ahmed A, McCunniff M, Overman P, Mathew M. Obesity and dental caries in children aged 2–6 years in the united states: National health and nutrition examination survey 1999–2002. *J Public Health Dent*. 2008; 68(4):227–233. [PubMed: 18384534]
58. Bailleul-Forestier I, Lopes K, Souames M, Azoguy-Levy S, Frelut ML, Boy-Lefevre ML. Caries experience in a severely obese adolescent population. *Int J Paediatr Dent*. 2007; 17(5):358–363. [PubMed: 17683325]
59. Hooley M, Skouteris H, Millar L. The relationship between childhood weight, dental caries and eating practices in children aged 4–8 years in australia, 2004–2008. *Pediatr Obes*. 2012

60. Marshall TA, Eichenberger-Gilmore JM, Broffitt BA, Warren JJ, Levy SM. Dental caries and childhood obesity: Roles of diet and socioeconomic status. *Community Dent Oral Epidemiol.* 2007; 35(6):449–458. [PubMed: 18039286]
61. McGuire J, Szabo A, Jackson S, Bradley TG, Okunseri C. Erosive tooth wear among children in the united states: Relationship to race/ethnicity and obesity. *Int J Paediatr Dent.* 2009; 19(2):91–98. [PubMed: 19250393]
62. Trasande L, Chatterjee S. The impact of obesity on health service utilization and costs in childhood. *Obesity (Silver Spring).* 2009; 17(9):1749–1754. [PubMed: 19300433]
63. Woolford SJ, Gebremariam A, Clark SJ, Davis MM. Persistent gap of incremental charges for obesity as a secondary diagnosis in common pediatric hospitalizations. *J Hosp Med.* 2009; 4(3): 149–156. [PubMed: 19301381]
64. Woolford SJ, Gebremariam A, Clark SJ, Davis MM. Incremental hospital charges associated with obesity as a secondary diagnosis in children. *Obesity (Silver Spring).* 2007; 15(7):1895–1901. [PubMed: 17636109]
65. El-Metainy S, Ghoneim T, Aridae E, Abdel Wahab M. Incidence of perioperative adverse events in obese children undergoing elective general surgery. *Br J Anaesth.* 2011; 106(3):359–363. [PubMed: 21149286]
66. Vu LT, Nobuhara KK, Lee H, Farmer DL. Determination of risk factors for deep venous thrombosis in hospitalized children. *J Pediatr Surg.* 2008; 43(6):1095–1099. [PubMed: 18558189]
67. Webb ML, Cerrato F, Rosen H, DiVasta AD, Greene AK, Labow BI. The effect of obesity on early outcomes in adolescents undergoing reduction mammoplasty. *Ann Plast Surg.* 2012; 68(3):257–260. [PubMed: 21629095]
68. Olutoye OA, Watcha MF, Andropoulos DB. Pediatric obesity: Observed impact in the ambulatory surgery setting. *J Natl Med Assoc.* 2011; 103(1):27–30. [PubMed: 21329243]
69. Adibi A, Kelishadi R, Beihaghi A, Salehi H, Talaei M. Sonographic fatty liver in overweight and obese children, a cross sectional study in isfahan. *Endokrynol Pol.* 2009; 60(1):14–19. [PubMed: 19224500]
70. Eminoglu TF, Camurdan OM, Oktar SO, Bideci A, Dalgic B. Factors related to non-alcoholic fatty liver disease in obese children. *Turk J Gastroenterol.* 2008; 19(2):85–91. [PubMed: 19110662]
71. Damaso AR, do Prado WL, de Piano A, et al. Relationship between nonalcoholic fatty liver disease prevalence and visceral fat in obese adolescents. *Dig Liver Dis.* 2008; 40(2):132–139. [PubMed: 18082476]
72. Suano de Souza FI, Silverio Amancio OM, Saccardo Sarni RO, et al. Non-alcoholic fatty liver disease in overweight children and its relationship with retinol serum levels. *Int J Vitam Nutr Res.* 2008; 78(1):27–32. [PubMed: 18654951]
73. Stordal K, Johannesdottir GB, Bentsen BS, Carlsen KC, Sandvik L. Asthma and overweight are associated with symptoms of gastro-oesophageal reflux. *Acta Paediatr.* 2006; 95(10):1197–1201. [PubMed: 16982489]
74. Teitelbaum JE, Sinha P, Micale M, Yeung S, Jaeger J. Obesity is related to multiple functional abdominal diseases. *J Pediatr.* 2009; 154(3):444–446. [PubMed: 19874760]
75. Wijga AH, Scholtens S, Bemelmans WJ, et al. Comorbidities of obesity in school children: A cross-sectional study in the PIAMA birth cohort. *BMC Public Health.* 2010; 10:184. [PubMed: 20380692]
76. Wang Y. Is obesity associated with early sexual maturation? A comparison of the association in american boys versus girls. *Pediatrics.* 2002; 110(5):903–910. [PubMed: 12415028]
77. Tussing-Humphreys LM, Liang H, Nemeth E, Freels S, Braunschweig CA. Excess adiposity, inflammation, and iron-deficiency in female adolescents. *J Am Diet Assoc.* 2009; 109(2):297–302. [PubMed: 19167957]
78. Samartzis D, Karppinen J, Mok F, Fong DY, Luk KD, Cheung KM. A population-based study of juvenile disc degeneration and its association with overweight and obesity, low back pain, and diminished functional status. *J Bone Joint Surg Am.* 2011; 93(7):662–670. [PubMed: 21471420]
79. Pfeiffer M, Kotz R, Ledl T, Hauser G, Sluga M. Prevalence of flat foot in preschool-aged children. *Pediatrics.* 2006; 118(2):634–639. [PubMed: 16882817]

80. Krul M, van der Wouden JC, Schellevis FG, van Suijlekom-Smit LW, Koes BW. Musculoskeletal problems in overweight and obese children. *Ann Fam Med*. 2009; 7(4):352–356. [PubMed: 19597173]
81. Hacıhamdioglu B, Okutan V, Yozgat Y, et al. Abdominal obesity is an independent risk factor for increased carotid intima-media thickness in obese children. *Turk J Pediatr*. 2011; 53(1):48–54. [PubMed: 21534339]
82. Giannini C, de Giorgis T, Scarinci A, et al. Obese related effects of inflammatory markers and insulin resistance on increased carotid intima media thickness in pre-pubertal children. *Atherosclerosis*. 2008; 197(1):448–456. [PubMed: 17681348]
83. Bhattacharjee R, Kim J, Alotaibi WH, Kheirandish-Gozal L, Capdevila OS, Gozal D. Endothelial dysfunction in children without hypertension: Potential contributions of obesity and obstructive sleep apnea. *Chest*. 2012; 141(3):682–691. [PubMed: 22030801]
84. Schaeffer AJ, Feng Z, Trock BJ, et al. Medical comorbidities associated with pediatric kidney stone disease. *Urology*. 2011; 77(1):195–199. [PubMed: 20970831]
85. Agranat-Meged AN, Deitcher C, Goldzweig G, Leibenson L, Stein M, Galili-Weisstub E. Childhood obesity and attention deficit/hyperactivity disorder: A newly described comorbidity in obese hospitalized children. *Int J Eat Disord*. 2005; 37(4):357–359. [PubMed: 15856493]
86. Kim J, Mutyala B, Agiovlasis S, Fernhall B. Health behaviors and obesity among US children with attention deficit hyperactivity disorder by gender and medication use. *Prev Med*. 2011; 52(3–4):218–222. [PubMed: 21241728]
87. Erhart M, Herpertz-Dahlmann B, Wille N, Sawitzky-Rose B, Holling H, Ravens-Sieberer U. Examining the relationship between attention-deficit/hyperactivity disorder and overweight in children and adolescents. *Eur Child Adolesc Psychiatry*. 2012; 21(1):39–49. [PubMed: 22120761]
88. Rojo L, Ruiz E, Dominguez JA, Calaf M, Livianos L. Comorbidity between obesity and attention deficit/hyperactivity disorder: Population study with 13–15-year-olds. *Int J Eat Disord*. 2006; 39(6):519–522. [PubMed: 16609984]
89. Marks S, Shaikh U, Hilty DM, Cole S. Weight status of children and adolescents in a telepsychiatry clinic. *Telemed J E Health*. 2009; 15(10):970–974. [PubMed: 20028189]
90. Eschenbeck H, Kohlmann CW, Dudey S, Schurholz T. Physician-diagnosed obesity in German 6- to 14-year-olds. prevalence and comorbidity of internalising disorders, externalising disorders, and sleep disorders. *Obes Facts*. 2009; 2(2):67–73. [PubMed: 20054208]
91. Fiese BH, Everhart RS, Wildenger L. Wheezing, sleeping, and worrying: The hidden risks of asthma and obesity in school-age children. *Psychology in the Schools*. 2009; 46(8):728–738.
92. Hillman JB, Dorn LD, Bin H. Association of anxiety and depressive symptoms and adiposity among adolescent females, using dual energy X-ray absorptiometry. *Clin Pediatr (Phila)*. 2010; 49(7):671–677. [PubMed: 20356924]
93. Gibson LY, Byrne SM, Blair E, Davis EA, Jacoby P, Zubrick SR. Clustering of psychosocial symptoms in overweight children. *Aust N Z J Psychiatry*. 2008; 42(2):118–125. [PubMed: 18197506]
94. Datar A, Sturm R. Childhood overweight and parent- and teacher-reported behavior problems: Evidence from a prospective study of kindergartners. *Arch Pediatr Adolesc Med*. 2004; 158(8):804–810. [PubMed: 15289255]
95. Anderson SE, Cohen P, Naumova EN, Must A. Association of depression and anxiety disorders with weight change in a prospective community-based study of children followed up into adulthood. *Arch Pediatr Adolesc Med*. 2006; 160(3):285–291. [PubMed: 16520448]
96. Bell LM, Byrne S, Thompson A, et al. Increasing body mass index z-score is continuously associated with complications of overweight in children, even in the healthy weight range. *J Clin Endocrinol Metab*. 2007; 92(2):517–522. [PubMed: 17105842]
97. Vila G, Zipper E, Dabbas M, et al. Mental disorders in obese children and adolescents. *Psychosom Med*. 2004; 66(3):387–394. [PubMed: 15184702]
98. Merikangas AK, Mendola P, Pastor PN, Reuben CA, Cleary SD. The association between major depressive disorder and obesity in US adolescents: Results from the 2001–2004 national health and nutrition examination survey. *J Behav Med*. 2012; 35(2):149–154. [PubMed: 21479835]

99. Bjornelv S, Nordahl HM, Holmen TL. Psychological factors and weight problems in adolescents. the role of eating problems, emotional problems, and personality traits: The young-HUNT study. *Soc Psychiatry Psychiatr Epidemiol*. 2011; 46(5):353–362. [PubMed: 20238097]
100. Huang L, Tao FB, Wan YH, et al. Self-reported weight status rather than BMI may be closely related to psychopathological symptoms among mainland chinese adolescents. *J Trop Pediatr*. 2011; 57(4):307–311. [PubMed: 19797398]
101. Lumeng JC, Gannon K, Cabral HJ, Frank DA, Zuckerman B. Association between clinically meaningful behavior problems and overweight in children. *Pediatrics*. 2003; 112(5):1138–1145. [PubMed: 14595059]
102. Hainsworth KR, Davies WH, Khan KA, Weisman SJ. Co-occurring chronic pain and obesity in children and adolescents: The impact on health-related quality of life. *Clin J Pain*. 2009; 25(8): 715–721. [PubMed: 19920723]
103. Nadeau K, Kolotkin RL, Boex R, et al. Health-related quality of life in adolescents with comorbidities related to obesity. *J Adolesc Health*. 2011; 49(1):90–92. [PubMed: 21700164]
104. de Beer M, Hofsteenge GH, Koot HM, Hirasings RA, Delemarre-van de Waal HA, Gemke RJ. Health-related-quality-of-life in obese adolescents is decreased and inversely related to BMI. *Acta Paediatr*. 2007; 96(5):710–714. [PubMed: 17381471]
105. Al-Akour NA, Khader YS, Khassawneh MY, Bawadi H. Health-related quality of life of adolescents with overweight or obesity in the north of jordan. *Child Care Health Dev*. 2012; 38(2):237–243. [PubMed: 21615771]
106. Schwimmer JB, Burwinkle TM, Varni JW. Health-related quality of life of severely obese children and adolescents. *JAMA*. 2003; 289(14):1813–1819. [PubMed: 12684360]
107. Kuhl ES, Rausch JR, Varni JW, Stark LJ. Impaired health-related quality of life in preschoolers with obesity. *J Pediatr Psychol*. 2012; 37(10):1148–1156. [PubMed: 22976509]
108. Bayer O, Rosario AS, Wabitsch M, von Kries R. Sleep duration and obesity in children: Is the association dependent on age and choice of the outcome parameter? *Sleep*. 2009; 32(9)
109. Nixon GM, Thompson JM, Han DY, et al. Short sleep duration in middle childhood: Risk factors and consequences. *Sleep*. 2008; 31(1):71–78. [PubMed: 18220080]
110. Liu X, Forbes EE, Ryan ND, Rofey D, Hannon TS, Dahl RE. Rapid eye movement sleep in relation to overweight in children and adolescents. *Arch Gen Psychiatry*. 2008; 65(8):924–932. [PubMed: 18678797]
111. Kohler M, Lushington K, Couper R, et al. Obesity and risk of sleep related upper airway obstruction in caucasian children. *J Clin Sleep Med*. 2008; 4(2):129–136. [PubMed: 18468311]
112. Carno MA, Modrak J, Short R, Ellis ER, Connolly HV. Sleep associated gas exchange abnormalities in children and adolescents with habitual snoring. *Pediatr Pulmonol*. 2009; 44(4): 364–372. [PubMed: 19283839]
113. Kohler MJ, Thormaehlen S, Kennedy JD, et al. Differences in the association between obesity and obstructive sleep apnea among children and adolescents. *J Clin Sleep Med*. 2009; 5(6):506–511. [PubMed: 20465015]
114. Wake M, Hardy P, Sawyer MG, Carlin JB. Comorbidities of overweight/obesity in australian preschoolers: A cross-sectional population study. *Arch Dis Child*. 2008; 93(6):502–507. [PubMed: 18218662]
115. Lee A, Ho MM, Keung VM. Global epidemics of childhood obesity is hitting a ‘less industrialized’ corner in asia: A case study in macao. *Int J Pediatr Obes*. 2011; 6(2–2):e252–6. [PubMed: 21649474]
116. Karatzi K, Protogerou A, Rarra V, Stergiou GS. Home and office blood pressure in children and adolescents: The role of obesity. the arsakeion school study. *J Hum Hypertens*. 2009; 23(8):512–520. [PubMed: 19129855]
117. Chu NF, Pan WH. Prevalence of obesity and its comorbidities among schoolchildren in taiwan. *Asia Pac J Clin Nutr*. 2007; 16 (Suppl 2):601–607. [PubMed: 17724001]
118. Bhowmick SK, Dasari G, Levens KL, Rettig KR. The prevalence of elevated serum thyroid-stimulating hormone in childhood/adolescent obesity and of autoimmune thyroid diseases in a subgroup. *J Natl Med Assoc*. 2007; 99(7):773–776. [PubMed: 17668643]

119. Fortmeier-Saucier L, Savrin C, Heinzer M, Hudak C. BMI and lipid levels in mexican american children diagnosed with type 2 diabetes. *Worldviews Evid Based Nurs.* 2008; 5(3):142–147. [PubMed: 19076913]
120. Nguyen S, McCulloch C, Brakeman P, Portale A, Hsu CY. Being overweight modifies the association between cardiovascular risk factors and microalbuminuria in adolescents. *Pediatrics.* 2008; 121(1):37–45. [PubMed: 18166555]
121. CDC. [Accessed November, 2012] Overweight and obesity:Basics about childhood obesity. <http://www.cdc.gov/obesity/childhood/basics.html>. Updated 2012
122. Physical status: The use and interpretation of anthropometry. report of a WHO expert committee. *World Health Organ Tech Rep Ser.* 1995; 854:1–452. [PubMed: 8594834]
123. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: International survey. *BMJ.* 2000; 320(7244):1240–1243. [PubMed: 10797032]
124. Nigg C, Shor B, Tanaka CY, Hayes DK. Adolescent at-risk weight (overweight and obesity) prevalence in hawai'i. *Hawaii Med J.* 2011; 70(7 Suppl 1):4–10.

Table 1

Psychological/Behavioral Co-morbidity	Age range	Ethnicity	Author, Year
<i>Internalizing Disorders</i>			
Anxiety and depression	9–18 years	92% white; 8% other	Anderson, 2006
Anxiety and depression	11–17 years	62% white; 38% non-white	Hillman, 2010
Depression	12–19 years	16% non-Hispanic white, 21% non-Hispanic black, 18.8% Mexican-Americans	Merikangas, 2012
General internalizing symptoms	5–12 years	55% White; 29% African American; 3% Hispanic; 1% Native American; 12% mixed race	Fiese, 2009
<i>Externalizing Disorders</i>			
ADHD	6–17 years	***62(62)% white; 15(14)% AA; 16(17)% Hispanic; 7(7)% other	Kim, 2011
ADHD	6–18 years	85% Hispanic	Mirza, 2004
Behavioral problems	* 4–5 years	** Girls 56 (65)% white ** Boys 66 (59) % white	Datar, 2004
Behavioral problems	8–11 years	53% white; 27% black; 20% Hispanic	Lumeng, 2003
Externalizing symptoms	5–12 years	55% White; 29% African American; 3% Hispanic; 1% Native American; 12% mixed race	Fiese, 2009
<i>Sleep Problems</i>			
Obstructive sleep apnea syndrome	2–17 years	---	Carno, 2009
Obstructive sleep apnea (OSA)	3–5 years; 13–16 years	***33(38)% White non-Hispanic; 37(46)% black non-Hispanic, 27(5)% Hispanic; 3(3)% other; 3(9)% American Indian	Lazorick, 2011
Obstructive sleep apnea, daytime somnolence	6–18 years	85% Hispanic	Mirza, 2004
Sleep disruptions	5–12 years	55% White; 29% African American; 3% Hispanic; 1% Native American; 12% mixed race	Fiese, 2009
Sleep problems	7–17 years	85% White; 12% Black; 3% other	Liu, 2008
Snoring	6–18 years	85% Hispanic	Mirza, 2004
<i>Other</i>			
Disordered eating	14–18 years	83% African-American; 7% white; 2% Hispanic; 2% other; 6% unknown	Trent, 2009
Psychological Conditions	<21 years	---	Marks, 2009
Medical Co-morbidity	Age range	Ethnicity	Author, Year
<i>Metabolic Risk Factors</i>			
Abnormal cholesterol, HDL, and triglyceride levels	8–20 years	100% Mexican-Americans	Fortmeier-Saucier, 2008
Hypertension	3–5 years; 13–16 years	***33(38)% White non-Hispanic; 37(46)% black non-Hispanic, 27(5)% Hispanic; 3(3)% other; 3(9)% American Indian	Lazorick, 2011

Psychological/Behavioral Co-morbidity	Age range	Ethnicity	Author, Year
Cardiovascular risk profile High Blood pressure	6–18 years	85% Hispanic	Mirza, 2004
High blood pressure	8–17 years	Mexican-American; Puerto-Rican; Cuban-American	Din-Dzietham, 2007
High blood pressure	14–18 years	83% African-American; 7% white; 2% Hispanic; 2% other; 6% unknown	Trent, 2009
Gastrointestinal complaints	2–20 years	---	Teitelbaum, 2009
Hypertension (HTN); left ventricular hypertrophy (LVH).	13–19 years	70% Caucasian; 9% African-American; 14% Hispanic; 6% other	Movahed, 2011
Metabolic syndrome	12–18 years	74% Latin; 26% Caribbean-black	Messiah, 2010
Metabolic syndrome	5–8 years	93% Hispanic; 7% other	Patino-Fernandez, 2008
Microalbuminuria and cardiovascular risk factors	12–19 years	27% non-Hispanic white; 31% non-Hispanic black; 39% Hispanic; 3% other	Nguyen, 2008
Thyroid function	6–17 years	---	Bhowmick, 2007
Pre-diabetes	3–5 years; 13–16 years	*** 33(38)% White non-Hispanic; 37(46)% black non-Hispanic, 27(5)% Hispanic; 3(3)% other; 3(9)% American Indian	Lazorick, 2011
<i>Asthma</i>			
Asthma	12–19 years	60% Caucasian; 19% African-American; 12% Hispanic; 9% biracial/other	Abramson, 2008
Asthma	5–12 years	55% White; 29% African American; 3% Hispanic; 1% Native American; 12% mixed race	Fiese, 2009
Asthma	7–18 years	*** 57(60)% White; 6(4)% Black; 29(28)% Hispanic; 8(9)% other	Gilliland, 2003
Asthma	3–5 years; 13–16 years	*** 33(38)% White non-Hispanic; 37(46)% black non-Hispanic, 27(5)% Hispanic; 3(3)% other; 3(9)% American Indian	Lazorick, 2011
Asthma	6–18 years	85% Hispanic; 15% other	Mirza, 2004
Asthma	9–22 years	96% Native American; 4% other	Noonan, 2010
Asthma	14–18 years	83% African-American; 7% white; 2% Hispanic; 2% other; 6% unknown	Trent, 2009
<i>Dental Health</i>			
Dental cavities	2–6 years	61% non-Hispanic White; 14% non-Hispanic Black; 20% Hispanic; 6% other	Hong, 2008
Erosive tooth wear	13–19 years	65% white; 15% African-American; 16% Hispanic; 4% other	McGuire, 2008
<i>Medical Expenditures</i>			
Health care expenditures	6–19 years	15% non-Hispanic Black; 18% Hispanic; 4% Asian; 63% other/unspecified	Trasande, 2009
<i>Other</i>			
Back pain, gastroesophageal reflux disease, acanthosis nigricans	3–5 years; 13–16 years	*** 33(38)% White non-Hispanic; 37(46)% black non-Hispanic, 27(5)% Hispanic; 3(3)% other; 3(9)% American Indian	Lazorick, 2011
Iron deficiency	12–17 years	** 27 (32)% non-Hispanic white; 40 (30)% non-Hispanic African-American; 33(39)% Hispanic	Tussing-Humphreys, 2009

Psychological/Behavioral Co-morbidity	Age range	Ethnicity	Author, Year
Complications after reduction mammoplasty surgery	13–20 years	---	Webb, 2012
Deep venous thrombosis	1–17 years	---	Vu, 2008
Endothelial dysfunction	4–12 years	59% white; 31% black; 5% Hispanic; 5% other	Bhattacharjee, 2012
Sexual maturation	8–14 years	26(25)% non-Hispanic white; 35(36)% non-Hispanic black; 35(35)% Mexican-American; 4(4)% other	Wang, 2002
Surgery	3–17 year	35% Caucasian; 17% African-American; 42% Hispanic; 11% other	Olutoye, 2011

---No ethnicity data available

* Estimated range based on mean and standard deviation

** Overweight/obese(healthy weight)%

*** Age at time 1(Age at time 2)%

**** Girls (boys)%

Table 2

Co-morbidities of Childhood Obesity from International Studies

Psychological/Behavioral comorbidity	Age range		Author, Year
<i>Internalizing Disorders</i>			
Anxiety and depression	6–13 years	Australia	Bell, 2007
Anxiety and depression	6–13 years	Australia	Bell, 2011
Anxiety Depression Self-esteem	13–19 years	Norway	Bjornelv, 2011
Depression, self-esteem	8–13 years	Western Australian	Gibson, 2008
Internalizing disorders	6–14 years	German	Eschenbeck, 2009
Internalizing disorders	5–17 years	France	Vila, 2004
<i>Externalizing Disorders</i>			
ADHD	8–17 years	Israel	Agranat-Meged, 2005
ADHD	11–17 years	Germany	Erhart, 2012
ADHD	13–15 years	Spain	Rojo, 2006
Externalizing disorders	6–14 years	German	Eschenbeck, 2009
Externalizing disorders	5–17 years	France	Vila, 2004
<i>Sleep Problems</i>			
Obstructive sleep apnea (OSA)	6–13 years	Australia	Bell, 2007
Obstructive sleep apnea syndrome (OSAS)	2–18 years	Australia	Kohler, 2009
Sleep disorders	6–14 years	German	Eschenbeck, 2009
Sleep duration	3–10 years	Germany	Bayer, 2009
Sleep duration	7 years	New Zealand	Nixon, 2008
Sleep problems	*5–11 years	Australia	Wake, 2010
<i>Other</i>			
Body dissatisfaction, eating disorder symptoms, bullying	8–13 years	Western Australian	Gibson, 2008
Bullying	6–13 years	Australia	Bell, 2011
Dieting behaviors	*5–11 years	Australia	Wake, 2010
Psychopathological symptoms health not associated	12–24 years	China	Huang, 2011
Social skills	5–17 years	France	Vila, 2004
Medical comorbidities			
<i>Metabolic Risk Factors</i>			
Blood pressure	6–13 years	Taiwan	Chu, 2007
Blood pressure, HDL-cholesterol, triglycerides and carbohydrate metabolism	1–20 years	Europe	I'Allemand, 2008
Blood pressure	6–18 years	Greece	Karatzis, 2009
Blood pressure	10–18 years	Poland	Krzyzaniak, 2011
Cardiovascular risk factors	3–18 years	The Netherlands	Langens, 2008
Cardiovascular risk factors	8–15 years	Portugal	Ribeiro, 2004
Cardiovascular risk factors and metabolic syndrome	6–16 years	Turkey	Serap, 2007
Carotid intima-media thickness increased	*7–12 years	Turkey	Hacıhamdioglu, 2011
Gastro-oesophageal reflux	7–16 years	Norwegian	Stordal, 2006

Psychological/Behavioral comorbidity	Age range		Author, Year
Hypertension, hyperlipidemia	17 years	Israel	Levin, 2010
Hypertension	11–17 years	Ludhiana	Mohan, 2004
Hypertension, LDL, hypertriglyceridemia	7–8 years	Taiwan	Ng, 2004
Hypertension	5–11 years	Australia	Wake, 2010
Metabolic risk	1–18 years	Germany	Flechtner-Mors, 2012
Metabolic risk	6–18 years	China	Lee, 2011
Metabolic risk	6–18 years	Germany	Reinehr, 2010
Metabolic syndrome	1–17 years	Berlin	Dannemann, 2011
Metabolic syndrome	12–14 years	Mexico	Del-Rio-Navarro, 2010
Metabolic syndrome	12–18 years	Italy Germany	Lafortuna, 2010
Non-alcoholic fatty liver disease	15–19 years	Brazil	Dámaso, 2008
Non-alcoholic fatty liver disease (NAFLD)	7–13 years	Turkey	Emino lu, 2008
Non-alcoholic fatty liver disease	6–10 years	Brazil	Suano de Souza, 2008
Sonographic Fatty Liver	6–18 years	Iran	Adibi, 2009
Type 2 diabetes	17 years	Israel	Levin, 2010
<i>Asthma</i>			
Asthma	6–11 years	Greece	Spathopoulos, 2009
Asthma	13–14 years	Macedonia	Vlaski, 2006
Asthma symptoms	6–15 years	France	Mahut, 2012
Asthma	4–5 years	Australia	Tai, 2009
Asthma Atopic disease	5–16 years	Portugal	Silva, 2007
Asthma	2–16 years	Italy	Vignolo, 2005
Asthma	5–11 years	Australia	Wake, 2010
Asthma	11–12 years	New Zealand	Wickens, 2005
<i>Dental health</i>			
Cavities	6–10 years	Germany	Willershausen, 2007
<i>Surgery</i>			
Perioperative implications	2–16 years	Egypt	El-Metainy, 2011
<i>Other</i>			
Airway hyper-responsiveness and exercise-induced bronchospasm frequencies	9–15 years	Turkey	Ulger, 2006
Airway hyper-responsiveness	10–12 years	Korea	Jang, 2006
Allergy	12–17 years	Vietnam	Irie, 2005
Atopy	6–11 years	Greece	Spathopoulos, 2009
Disc degeneration	13–20 years	China	Samartzis, 2011
Flat foot	3–6 years	Austria	Pfeiffer, 2006
Impaired glucose tolerance (IGT) hyperinsulinism dyslipidaemia raised alanine transaminase	6–13 years	Australia	Bell, 2011
Intima media thickness (IMT)	*6–10 years	Italy	Giannini, 2008
Muscle and joint problems	5–11 years	Australia	Wake, 2010

Psychological/Behavioral comorbidity	Age range		Author, Year
Musculoskeletal pain, headaches, acanthosis nigricans	6–13 years	Australia	Bell, 2007
Musculoskeletal pain	6–13 years	Australia	Bell, 2011
Musculoskeletal problems	2–17 years	The Netherlands	Krul, 2009
Poor health status School absenteeism	8 years	The Netherlands	Wijga, 2010
Risk factors for atherosclerosis	6–20 years	Poland	Glowinska, 2003
Upper Airway Obstruction	4–12 years	Australia	Kohler, 2008